

# INCL Intra-nuclear Cascade and ABLA De-excitation Models in Geant4

Pekka KAITANIEMI<sup>1,3\*</sup>, Alain BOUDARD<sup>1</sup>, Sylvie LERAY<sup>1</sup>, Joseph CUGNON<sup>2</sup>, Davide MANCUSI<sup>2</sup>  
On behalf of the Geant4 collaboration

<sup>1</sup> CEA/Saclay, DSM/IRFU/SPhN, 91191 Gif-sur-Yvette Cedex, France

<sup>2</sup> University of Liège, Allée du 6 Août 17, bât B5, B-4000 Liège, Belgium

<sup>3</sup> Helsinki Institute of Physics, P.O.Box 64 (Gustaf Hällström street 2), FIN-00014 University of Helsinki, Finland

We present INCL intra-nuclear cascade and ABLA de-excitation (evaporation/fission) models that can be used to calculate collisions between projectile particles and nuclei at energies from 0.2 to 3 GeV. Supported projectiles are proton, neutron, pions and light ions up to alpha. The optimal target materials range from carbon to uranium.

We have extended the INCL model light ion projectile support up to carbon-ions and compared the results against thin and thick target data and Geant4 Binary cascade model. Both thin and first thick target results of the INCL model are in good agreement with experimental data.

**KEYWORDS:** *Geant4, intra-nuclear cascade, Monte Carlo, ion-ion collisions*

## I. Introduction

INCL<sup>1)</sup> intra-nuclear cascade model calculates the reactions between nucleon, pion or light ion projectiles with target nuclei. After the cascade an excited remnant nucleus is de-excited using the ABLA<sup>2-4)</sup> evaporation/fission code. INCL/ABLA alone is a thin-target simulation code. In order to perform calculations that involve complex geometries and multiple composite materials INCL must be embedded into a transport code that can offer this functionality. Two such transport codes are Geant4<sup>5)</sup> and MCNPX<sup>6)</sup>. The original FORTRAN version of INCL4.2 has been included in MCNPX<sup>7)</sup>. The same version of INCL was later translated to C++ and included in Geant4<sup>8)</sup>.

The combination of INCL/ABLA with transport code has opened new possibilities for practical applications. Some of the more traditional applications for intra-nuclear cascade models are nuclear waste transmutation and Accelerator Driven Systems (ADS) studies which mainly deal with heavy target materials and nucleon beams. In recent years there has also been an increasing interest in medical applications of these Monte Carlo codes. One such application is the treatment of tumors using light ions, especially carbon beams with energies of a few hundred MeV per nucleon. Another interesting application for ion-ion collisions are the spacecraft radiation damage and shielding studies. In this case the relevant ions range from protons to iron ions with energies up to a few GeV.

This has motivated us to extend the light ion projectiles supported by INCL. In the previous release of INCL4.2 we supported light ion projectiles up to alpha. In this new version we have extended this support, in a fairly crude way, up to

carbon. In this paper we present our first results and compare against Geant4 Binary cascade which offers similar support for light ion projectiles as INCL.

In the future we hope to further extend the ion-ion support of the INCL intra-nuclear cascade model and make it more realistic. This, and many technical shortcomings of the INCL codebase that have their roots in the FORTRAN77 legacy of the original code, have motivated us to perform a complete redesign of the INCL simulation code in object oriented C++. We will discuss some of our motivations and main ideas of the redesign work and how they will help us to develop our model further.

## II. INCL4.2 light ion extension

The INCL light ion extension consists of two main parts: handling of the projectile as a collection of individual nucleons and de-excitation of the projectile fragments after the reaction. The main cascade in the target nucleus is treated following the standard INCL cascade procedure as described in reference<sup>1)</sup>.

### 1. Projectile as a collection of nucleons

The projectile is modeled as a collection of independent nucleons with gaussian momentum and position distributions<sup>9)</sup>. For the position distribution of the projectile nucleons we use the realistic standard deviation of the projectile ion. The standard deviation of the momentum distribution is fixed to 100 MeV/c. When generating momentum distribution for the projectile nucleons we first generate it in the rest frame of the projectile and then boost it to the laboratory frame. The sum of the energies of the projectile nucleons is equal to the total energy of the projectile. The momentum of the projectile nu-

\*Corresponding Author, E-mail:pekka.kaitaniemi@gmail.com

cleons is slightly rescaled due to the fact that they are on mass shell ( $E^2 - p^2 = m^2$ ) which is not compatible with the conservation of energy and momentum due to the binding energy of the projectile nucleus.

Some projectile nucleons miss the target. These are considered *projectile spectators*. Some of the nucleons that enter the target nucleus can pass through it without any collisions at all and are also considered as projectile spectators. The rest of the projectile nucleons proceed to produce an intra-nuclear cascade in the target as described in reference<sup>1)</sup>.

At the end of the cascade all projectile spectators are combined into a *spectator nucleus*. The mass, charge and momentum of this nucleus are directly determined by summation. The excitation energy is determined as follows. In the initialization of the cascade, a list of ion-projectile Fermi momenta has been randomly chosen corresponding to individual kinetic energies of the projectile nucleons. The removal of some projectile nucleons by the interaction with the target nucleus is interpreted as holes for the the projectile spectator nucleus. Its excitation energy is simply the energy released in packing the spectator nucleons in the lowest energy states of the individual kinetic energy list.

## 2. De-excitation

After the cascade we need to de-excite both the projectile spectator nucleus and the remnant of the target nucleus. The spectator nucleus is always de-excited using the Geant4 Fermi break-up model. Fermi break-up is also used for the light cascade remnants (below mass 17) as well. The heavy remnants are treated with the ABLA fission/evaporation code. The ABLA version in Geant4 provides us with evaporation of protons, neutrons and alphas that competes with fission channel for sufficiently heavy remnant nuclei.

## III. Results

We have compared the new version of INCL against experimental data and the Geant4 Binary cascade. These calculations have been done using a test version of INCL and Geant4 version `geant4-09-03-ref-05` which is an internal Geant4 development release from May of 2010.

### 1. Thin target calculations

Double-differential energy spectra for neutrons produced in reaction  $C + C$  at 135 MeV/nucleon are shown in **Figure 1**. Agreement of the INCL4.2 model with experiment is very good in the forward region. INCL combined with Geant4 Fermi break-up reproduces the quasi-elastic peak very well, as can be seen in the energy spectrum for angle  $0^\circ$ . The agreement with experimental data is worse for angles above  $30^\circ$ .

In the case of  $C + C$  at 290 MeV/nucleon (**Figure 2**) INCL performance is still fairly good, but not quite perfect, in the forward region. Both models show fairly similar levels of agreement with experiment. Overall INCL seems to reproduce the shape of the distribution slightly better than Binary cascade.

We have also compared oxygen and chlorine fragmentation results of INCL4.2 (Geant4 C++ version), INCL4.3 (FORTRAN version) and Binary cascade against experimental data.

The results are shown in **Figure 3**. It should be noted that in the experiment carbon was used as target but for technical reasons all calculations with the models were done using inverse kinematics (i.e. carbon as the projectile) and therefore in the calculation results we actually look at target fragmentation. The results of these comparisons show that the combination of INCL4.2 and Geant4 Fermi break-up performs quite well when compared against Binary Cascade and the FORTRAN version of INCL4.3 that uses Fermi break-up taken from the LAHET code<sup>10)</sup>.

The oxygen and chlorine fragmentation plots in **Figure 3** show a slight disagreement between the FORTRAN version INCL4.3 and the C++ version of INCL4.2. This difference can presumably be mostly attributed to the main difference between INCL4.2 and INCL4.3: light cluster emission in the cascade stage. It should also be noted that the Fermi break-up models used in the calculations come from different sources. In the FORTRAN version INCL4.3 we use Fermi break-up implementation from the LAHET package and in INCL4.2 Geant4 version we use the Geant4 Fermi break-up model.

### 2. Thick water target

We have started doing preliminary tests using a full Geant4 simulation with thick water target. For this purpose we use a modified version of the Geant4 hadron therapy example application included in the standard Geant4 distribution.


In our simulation setup we shoot a 200 MeV/nucleon carbon beam to a 12.78 cm thick water target and record the outgoing particles behind the target. A preliminary neutron production comparison between INCL and Geant4 Binary cascade is shown in **Figure 4**.

The physics performance of both INCL (using physics list `QGSP_INCL_ABLA`) and Binary cascade (using physics list `QGSP_BIC_EMY`) is quite good as far as neutron production is concerned.

## IV. Ongoing development: INCL redesign in C++

The design of the INCL FORTRAN code has remained relatively stable for more than a decade while physics performance and modeling features of the code have been significantly improved. This has made the codebase monolithic and difficult to develop further. It has become apparent that the current design has reached the end of its life. Since the Geant4 version of INCL4.2 is basically only a minimal translation of FORTRAN code to C++, sort of "FORTRAN in C++ syntax", it inherits all the problems of the FORTRAN version. Additionally the monolithic nature of the code makes it difficult to utilize the full potential of C++ capabilities.

Another important problem in the current development model of INCL is that we have several "forked" versions of the code: the main line of development (version INCL4.6) and the MCNPX version (currently INCL4.2, INCL4.5<sup>16)</sup> interface under development) in FORTRAN and INCL4.2 translation in C++ for Geant4. Additionally the light ion extension that was first introduced in INCL4.3 has now been added to the Geant4 version of INCL and will be released in 2010. The MCNPX version of INCL is essentially the same code as the standalone version, except that it has been modified to work



**Fig. 1** Double-differential neutron energy spectra for C + C at 135 MeV/nucleon. INCL4.2 with Geant4 Fermi break-up and Geant4 Binary cascade are compared against experimental data from reference<sup>11</sup>).

inside of a transport code. The redesign project gives us an opportunity to unify all these different versions under a single source tree and allows us to deliver the same physics features consistently for Geant4, MCNPX and as a standalone thin-target calculation code without duplication of effort.

The redesign project gives us an opportunity to revisit the physics ideas of INCL and experiment with features that would be very difficult to implement in current versions of INCL. Some examples of things that are difficult with the current codebase are changing the way particles are tracked or collisions are detected and implementing more realistic handling two of nuclear potentials in ion-ion collisions.

## V. Conclusions

We have presented the INCL intra-nuclear cascade model implementation in Geant4 and the INCL light ion projectile extension. The first results are in good agreement with experiment and physics performance of INCL is competitive with Geant4 Binary cascade.

We are now in good position to perform thick target studies and comparisons of INCL against experimental data in the case of light-ion collisions. Our first results are quite encouraging. However, there are still some open questions with regards to the physics list we should use with INCL. An especially important question is what to do for low energy collisions. In the physics list we used in this study we use INCL for nucleon, pion and ion projectiles from energy 3 GeV/nucleon down to 0 MeV/nucleon. Unfortunately some of the basic assumptions of intra-nuclear cascade start breaking down when energies fall significantly below 100 MeV/nucleon. One avail-

able option is to investigate the use of other potentially more suitable Geant4 models for low energy collisions and compare their results against those given by INCL.


The validation of the light ion extension will continue and it will be included in the Geant4 9.4 release in December 2010.

## Acknowledgment

We are grateful to José-Manuel Quesada and Mike Kelsey (Geant4 collaboration) for their insightful comments and suggestions for improvements concerning this paper. This work has been partly funded by Helsinki Institute of Physics and Fortum Foundation.


## References

- 1) A. Boudard et al., *Intra-nuclear cascade model for a comprehensive description of spallation reaction data*, Phys. Rev. C66 (2002) 044615
- 2) J. Benlliure et al., *Calculated nuclide production yields in relativistic collisions of fissile nuclei*, Nuc. Phys. A628 (1998) 458
- 3) J. J. Gaimard et al., Nuc. Phys. A531 (1991) 709
- 4) A. R. Junghans et al., Nuc. Phys. A629 (1998) 635
- 5) Geant4 collaboration, *Geant4 - a simulation toolkit*, Nucl. Instr. and Meth. A, 506, 2003, 250–303
- 6) J.S. Hendrix et al., AIP Conf. Proc. 768 (2005) 1188
- 7) Denise B. Pelowitz, et al., *INCL4.2-ABLA in MCNPX: MCNPX User's Manual, Version 2.5.0*, LA-CP-05-0369 (2005)
- 8) A. Heikkinen, P. Kaitaniemi, and A. Boudard, *Implementation of INCL4 cascade and ABLA evaporation codes in Geant4*, Journal of Physics: Conference Series 119 (2008) 032024, doi:10.1088/1742-6596/119/3/032024




**Fig. 2** Double-differential neutron energy spectra for C + C at 290 MeV/nucleon. INCL and Geant4 Binary cascade are compared against experimental data from reference<sup>12)</sup>.

- 9) A. Boudard, J. Cugnon, P. Kaitaniemi, S. Leray, D. Mancusi, *Simulation of light ion collisions from Intra Nuclear Cascade (INCL-Fermi Breakup) relevant for medical irradiations and radioprotection* AP/IE-08
- 10) R. E. and H. Liechtenstein, Report LA-UR-89-3014, Los Alamos National Laboratory, 1989
- 11) H. Sato et al., Physical Review C 64 (2001) 034607
- 12) Y. Iwata Physical Review C 64 (2001) 034607
- 13) C. Zeitlin et al., <http://fragserver.lbl.gov>
- 14) C. Zeitlin et al., Physical Review C 77 (2008) 034605
- 15) K. Gunzert-Marx, H. Iwase, D. Schardt, and R.S. Simon, *Secondary beam fragments produced by 200 MeV/u <sup>12</sup>C ions in water and their dose contribution in radiotherapy*, New Journal of Physics 10 (2008), doi:10.1088/1367-2630/10/7/075003
- 16) J. Cugnon, A. Boudard, S. Leray, D. Mancusi, Proceedings of the International Topical Meeting on Nuclear Research Applications and Utilization of Accelerators, Vienna, May 4-8, 2009



**Fig. 3** Oxygen (left) and chlorine (right) fragmentation of INCL4.2 (C++) with Geant4 Fermi break-up for light fragments (mass 16 and below) and ABLA for heavier fragments, Binary cascade, and INCL4.3 (FORTRAN) compared against experimental data from references<sup>13, 14</sup>. For technical reasons all calculations are done using inverse kinematics.



**Fig. 4** Neutron double-differential energy spectra in C + water with target thickness of 12.78 cm at 200 MeV/nucleon. INCL4.2 with Geant4 Fermi break-up and Binary cascade are compared against data from reference<sup>15</sup>.